

Item #	Question
1	What areas are frequently missed during handwashing/sanitizing? (select best answer)
2	Application of an alcohol rub or gel reduces the number of bacteria on your hands more effectively than handwashing with plain or antimicrobial soap. (true/false)
3	When should you perform hand hygiene? (select all that apply)
4	When should soap and water be used (rather than alcohol rub)? (select all that apply)
5	Gloves are an appropriate substitute for hand hygiene. (true/false)
6	Staff can bring in their own hand sanitizer as long as it is for personal use. (true/false)
7	What is the main route of transmission of healthcare-associated pathogens between patients in a healthcare facility? (select best answer)
8	Which of the following should be avoided due to increased likelihood of hand colonization with healthcare-associated pathogens? (select best answer)
9	What is an indication for doffing gloves, performing hand hygiene and re-donning a new pair of gloves? (select best answer)
10	Alcohol-based hand rub contributes to antibiotic resistance. (true/false)
11	Hand hygiene should always be performed after doffing gloves, but is not always required before donning gloves. (true/false)
12	In addition to universal masking and eyewear, the following PPE should be worn when entering the room of a patient on Contact Precautions: (select best answer)
13	Patients on Contact Precautions may use the common areas of the unit (e.g. activity room or kitchen). (true/false)
14	When transporting a patient on Contact Precautions, which of the following are true? (select all that apply)
15	When caring for a patient on Contact Precautions, dedicate patient care equipment (e.g. thermometer, stethoscope) to the patient, or disinfect the equipment prior to using on another patient. (true/false)
16	In addition to universal masking and eyewear, the following PPE should be worn when entering the room of a patient on Enhanced Contact Precautions: (select best answer)
17	When caring for a patient on Enhanced Contact Precautions, dedicate patient care equipment (e.g. thermometer, stethoscope) to the patient, or disinfect the equipment prior to using on another patient. (true/false)
18	Patients on Enhanced Contact Precautions should remain in their room, only leaving for essential medical purposes. An ear-loop mask should be worn whenever outside of their room. (true/false)
19	When transporting a patient on Enhanced Contact Precautions, which of the following are true? (select all that apply)
20	In addition to universal masking and eyewear, the following PPE should be worn when entering the room of a patient on Droplet Precautions: (select best answer)
21	Patients on Droplet Precautions may use the common areas on the unit (e.g. the activity room or kitchen). (true/false)
22	When transporting a patient on Droplet Precautions, which of the following are true? (select all that apply)
23	In addition to universal masking and eyewear, the following PPE should be worn when entering the room of a patient on Contact + Droplet Precautions: (select best answer)
24	When caring for a patient on Contact + Droplet Precautions, dedicate patient care equipment (e.g. thermometer, stethoscope) to the patient, or disinfect the equipment prior to using on another patient. (true/false)

- 25 When transporting a patient on Contact + Droplet Precautions, which of the following are true? (select all that apply)
- 26 What PPE is required to enter the room of a patient on Airborne Precautions? (select best answers)
- 27 You should perform hand hygiene before AND after doffing your N95/PAPR. (true/false)
- 28 A patient on Airborne Precautions requires a negative pressure isolation room and the door must remain closed at all times. (true/false)
- 29 When entering the room of a patient on Airborne Precautions, it is ok if the outer door to the ante-room is still open when you open the inner door to the patient's room. (true/false)
- 30 All staff must wear a properly Fit-Tested N95 respirator and eye protection, or a PAPR, to enter an Airborne Precaution room. (true/false)
- 31 A patient on Airborne Precautions may leave their room as long as they wear an N95. (true/false)
- 32 When transporting a patient on Airborne Precautions, which of the following are true? (select all that apply)
- 33 What PPE should be worn when entering the room of a patient on Airborne + Contact Precautions: (select best answer)
- 34 A patient on Airborne + Contact Precautions can leave their room as long as they wear an N95, gown and gloves. (true/false)
- 35 What PPE should be worn when entering the room of a patient on Enhanced Contact + Droplet Precautions? (select best answer)
- 36 When caring for a patient on Enhanced Contact + Droplet Precautions, staff should: (select best answer)
- 37 When transporting a patient on Enhanced Contact + Droplet Precautions, which of the following are true? (select all that apply)
- 38 What PPE should be worn when entering the room of a patient on Continuous Aerosol-Generating Procedure Precautions? (select best answer)
- 39 What should you do before entering the room of a patient on Intermittent Aerosol-Generating Procedure Precautions? (select all that apply)
- 40 You are caring for a patient with cystic fibrosis who needs a nebulized medication. What is the appropriate PPE? (select all that apply)
- 41 You are caring for a patient with cystic fibrosis who needs a nebulized medication. What precaution signs would be displayed at the bedside? (select best answer)
- 42 You are suctioning the tracheostomy of a patient. What is the appropriate PPE? (select all that apply)
- 43 Please see the staff member entering Contact/Droplet Precaution room, and identify hand hygiene or PPE errors: (select all that apply)
- 44 Please see the staff member caring for a patient, and identify hand hygiene or PPE errors: (select all that apply)
- 45 Please see the staff member entering a precaution room, and identify hand hygiene or PPE errors: (select all that apply)
- 46 Please identify hand hygiene or PPE errors: (select all that apply)