Supplementary Table S1 Median values (25% - 75% quartile range) for Welfare Quality (WQ) measures in Year 0 as well as the respective change in Year 1 (difference Year 0 and Year 1) in Austrian dairy farms (n = 34)

| | | | Year 0 | | Difference Year 0 – Year 1 | |
|---|-----------------------|---|--------|-----------------------|----------------------------|-----------------------|
| WQ measures | Unit | WQ principle ¹ / criteria ² | Median | 25% – 75% quartile | Median | 25% – 75% quartile |
| Very lean cows | % | P1 / C1 | 2.9 | 0.0 - 6.2 | 0.0 | -3.1 – 2.9 |
| Duration of lying down movement | sec | P2 / C3 | 5.9 | 5.5 - 6.1 | -0.1 | -0.6 - 0.2 |
| Lying down movements with collisions | % | P2 / C3 | 33.3 | 12.5 - 42.9 | -9.3 | - 27.4 – 8.7 |
| Lying cows which lie partly/completely outside lying area | % | P2 / C3 | 4.2 | 0.0 - 13.9 | 0.1 | -3.7 - 2.1 |
| Cows with dirty lower legs | % | P2 / C3 | 42.9 | 27.1 - 57.2 | 4.9 | -22.2 – 16.2 |
| Cows with dirty udder | % | P2 / C3 | 62.9 | 54.5 - 75.7 | -17.5 | -28.73.3 |
| Cows with dirty flank and upper legs | % | P2 / C3 | 40.0 | 27.3 - 65.7 | -10.3 | -19.9 – 12.9 |
| Not lame cows | % | P3 / C6 | 60.6 | 50.0 - 74.3 | -3.4 | -10.5 - 4.9 |
| Moderately lame cows | % | P3 / C6 | 20.8 | 16.6 - 27.0 | 1.5 | -3.6 - 7.5 |
| Severely lame cows | % | P3 / C6 | 13.5 | 9.1 - 24.6 | 2.3 | -6.6 - 6.0 |
| Cows with no hairless patch and no lesion | % | P3 / C6 | 38.9 | 30.6 - 48.5 | 26.4 | 18.5 - 34.0 |
| Cows with at least one hairless patch and no lesion | % | P3 / C6 | 32.4 | 27.1 - 46.2 | -15.1 | -23.38.9 |
| Cows with at least one lesion | % | P3 / C6 | 20.0 | 12.1 - 33.3 | -10.6 | -18.2 - 0.9 |
| Frequency of coughing | per cow per 15 min | P3 / C7 | 0.1 | 0.1 - 0.2 | 0.0 | -0.1 – 0.0 |
| Cows with nasal discharge | % | P3 / C7 | 0.0 | 0.0 - 0.0 | 0.0 | 0.0 - 0.0 |
| Cows with ocular discharge | % | P3 / C7 | 0.0 | 0.0 - 3.2 | 0.0 | -2.9 - 0.0 |
| Cows with increased respiratory rate | % | P3 / C7 | 0.0 | 0.0 - 0.0 | 0.0 | 0.0 - 0.0 |

| Cows with diarrhoea | % | P3 / C7 | 6.9 | 3.8 - 18.3 | 5.9 | -6.3 – 10.3 |
|---|---------------------|----------|------|-------------|------|--------------------|
| Cows with vulvar discharge | % | P3 / C7 | 0.0 | 0.0 - 3.1 | 0.0 | 0.0 - 0.0 |
| Mastitis (milk somatic cell count > 400 000) | % | P3 / C7 | 7.2 | 3.2 - 12.3 | 1.9 | - 2.9 – 4.5 |
| Mortality during the last 12 months | % | P3 / C7 | 0.0 | 0.0 - 3.3 | 0.0 | -2.7 - 0.2 |
| Dystocia | % | P3 / C7 | 2.8 | 0.0 - 5.5 | 0.0 | -3.9 - 2.6 |
| Downer cows | % | P3 / C7 | 2.8 | 0.0 - 6.0 | 0-0 | -3.0 - 2.6 |
| Frequency of headbutts | per cow per hour | P4 / C9 | 0.7 | 0.4 - 1.2 | -0.1 | -0.5 – 0.2 |
| Frequency of other aggressive events (displacements, chasing, fighting, chasing-up) | per cow per hour | P4 / C9 | 0.3 | 0.2 - 0.7 | 0.0 | -0.2 - 0.3 |
| Cows that can be touched | % | P4 / C11 | 42.9 | 30.9 - 50.9 | -4.5 | -15.8 - 9.9 |
| Cows that can be approached by 50 cm but not touched | % | P4 / C11 | 41.4 | 34.7 - 52.2 | 9.1 | 1.9 - 17.4 |
| Cows that can be approached between 50 cm an 1 m | % | P4 / C11 | 10.3 | 5.7 - 19.3 | -2.9 | -8.8 - 0.0 |
| Cows that can't be approached | % | P4 / C11 | 0.0 | 0.0 - 4.3 | 0.0 | -3.9 - 0.0 |
| Positive emotional state | WQ-score | P4 / C12 | 25.6 | 15.8 - 37.8 | 0.8 | -16.3 – 9.8 |

¹ P: Welfare Quality principle; P1 Good feeding; P2 Good housing; P3 Good health; P4 Appropriate behaviour

² C: Welfare Quality criteria; C1 Absence of prolonged hunger; C3 Comfort around resting; C6 Absence of injuries; C7 Absence of disease; C9 Expression of social behaviours;

C11 Good human-animal relationship; C12 Positive emotional state