

**Supplemental Table 6. Isotopic Results from this Study. The Mattocks, Old Town, and Mimbres averages do not include MT07 and OT02 because they had C:N ratios outside the acceptable range of reasonably well-preserved collagen.**

Sample ID	$\delta^{15}\text{N}_{\text{co}}$ (‰ vs. AIR)	$\delta^{13}\text{C}_{\text{co}}$ (‰ vs. VPDB)	wt %N	wt %C	Atomic C:N
ER01	9.9	-9.3	16.0	43.3	3.2
ER02	8.8	-8.1	15.4	42.0	3.2
ER03	8.2	-8.5	15.7	42.7	3.2
ER04	6.9	-15.3	15.3	41.8	3.2
ER05	7.1	-7.4	14.9	40.5	3.2
ER06	9.3	-8.3	15.4	41.9	3.2
ER07	8.5	-8.9	15.6	42.5	3.2
ER08	8.9	-8.7	15.4	41.8	3.2
ER09	8.3	-9.9	15.5	42.4	3.2
ER10	9.7	-8.3	14.9	40.6	3.2
TU949	8.5	-6.9	15.5	38.8	2.9
TU950	8.6	-7.0	16.8	41.9	2.9
TU951	8.5	-7.1	16.1	40.1	2.9
TU952	8.6	-7.0	16.1	40.1	2.9
TU953	8.3	-7.0	15.8	39.5	2.9
TU954	8.6	-7.3	14.9	37.2	2.9
TU955	8.4	-8.5	15.5	39.0	2.9
<i>Elk Ridge avg.</i>	8.5	-6.8			
LR01	6.3	-8.0	14.9	41.2	3.2
LR02	6.3	-9.4	14.4	39.8	3.2
LR03	3.6	-16.9	14.3	39.7	3.2
<i>Lake Roberts avg.</i>	5.4	-11.4			
MT01	6.0	-16.8	14.5	40.1	3.2
MT02	7.7	-7.6	13.3	37.1	3.3
MT03	6.5	-7.7	14.6	40.1	3.2
MT04	10.2	-9.0	14.6	41.1	3.3
MT05	7.7	-7.5	14.5	40.4	3.3
MT06	6.0	-16.1	13.8	39.2	3.3
MT07	8.0	-7.1	10.5	38.5	4.3
<i>Mattocks Avg.</i>	7.4	-10.8			
OT01	7.0	-11.6	14.0	40.9	3.4
OT02	6.2	-24.1	1.3	9.6	8.5
<i>Old Town avg.</i>	7.0	-11.6			
WS01	6.5	-8.0	14.5	40.4	3.3
WS02	8.0	-11.0	14.2	40.2	3.3
<i>Wheaton-Smith avg.</i>	7.3	-9.5			
<i>Mimbres avg.</i>	7.8	-9.4			