

## Hautaniemi et al., Supplemental Digital Content 1

Regular medications used by the subjects: number of subjects with each type of medication.

Topical corticosteroids comprise both nasal and inhaled medications.

	<b>LTP5+PSe</b>	<b>LTP25+PSe</b>	<b>Placebo</b>	<b>All</b>
	<b>(n=36)</b>	<b>(n=35)</b>	<b>(n=33)</b>	<b>(n=104)</b>
Acetylsalicylic acid	0	2	0	2
Alfuzosin	1	0	0	1
Antiepileptic agents	0	2	0	2
Antimicrobials	1	0	0	1
Benzodiazepines	2	0	0	2
Brimonidine eye drops	0	0	1	1
Corticosteroids, topical	2	0	1	3
Dopamine agonists	0	1	0	1
Hormone replacement therapy	6	1	1	8
Flecainide	0	0	1	1
Glucosamine	0	1	0	1
Latanoprost eye drops	0	0	1	1
NSAID	0	0	1	1
Tizanidine	1	0	1	2
Proton pump inhibitor	1	0	0	1
Salazosulphapyridine	0	0	1	1
Salmeterol	1	0	0	1
SSRI	2	4	2	8
SNRI	2	0	0	2
Thyroxine	4	2	2	8
Timolol eye drops	0	0	1	1
Tramadol	1	0	0	1
Tricyclic antidepressant	1	0	0	1

NSAID, non-steroidal anti-inflammatory agent; SSRI, selective serotonin reuptake inhibitor; SNRI, serotonin-norepinephrine reuptake inhibitor