Bateman A, Fonagy P. Impact of clinical severity on outcomes of mentalisation-based treatment for borderline personality disorder. Br J Psychiatry doi: 10.1192/bjp.bp.112.121129

Data supplement DS1: Sample characteristics by severity criteria at baseline.

Severity indicators are dichotomized using median splits for the purposes of illustration except Number of BPD criteria, which was used as a binary variable in the analysis.

	Total sample ple N 134		Number of Axis I diagnoses				Number of BPD criteria				Number of Axis II diagnoses (incl. BPD)				Symptom distress (GSI)			
			2 or fewer 58		3 or more 76		5 criteria 81		6 or more 49		2 or fewer 77		3 or more 57		GSI < 2.01 67		GSI ≥ 2.01 67	
Sample N																		
Mean age (SD)	31.1	(7.7)	30.1	(7.0)	31.9	(8.2)	30.5	(7.9)	32.5	(7.4)	31.0	(8.0)	31.3	(7.6)	31.7	(7.6)	30.4	(7.8)
Male: n (%)	27	(20%)	12	(21%)	15	(20%)	13	(16%)	14	(29%)	9	(12%)	18	(32%)***	13	(19%)	14	(21%)
GAF: Mean (SD)	41.0	(8.4)	42.5	(8.5)	39.8	(8.2)	40.7	(9.0)	40.8	(7.3)	42.5	(8.2)	39.0	(8.3)**	41.7	(8.9)	40.3	(7.9)
Number Axis I: Mean (SD)	2.8	(1.3)	1.6	(0.6)	3.8	(0.8)***	2.7	(1.2)	1.1	(1.3)*	2.6	(1.3)	3.0	(1.3)*	2.7	(1.3)	2.9	(1.3)
GSI: Mean (SD)	2.0	(0.6)	1.9	(0.6)	2.0	(0.7)	2.0	(0.6)	1.2	(0.7)	1.9	(0.7)	2.1	(0.6)*	1.5	(0.4)	2.5	(2.4)***
BDI: Mean (SD)	29.5	(9.5)	28.5	(9.4)	30.3	(9.5)	29.3	(9.4)	30.5	(9.4)	28.5	(9.9)	30.9	(8.8)	24.7	(8.4)	34.2	(8.0)***
IIP: Mean (SD)	2.0	(0.5)	1.9	(0.5)	2.1	(0.5)	2.0	(0.5)	2.0	(0.5)	1.9	(0.5)	2.1	(0.5)*	1.8	(1.7)	2.3	(2.1)***
SAS: Mean (SD)	2.7	(0.6)	2.7	(0.5)	2.8	(0.6)	2.7	(0.5)	2.7	(0.6)	2.7	(0.6)	2.8	(0.5)	2.5	(2.4)	2.9	(2.7)**
Antisocial PD n (%)	37	(28%)	14	(24%)	23	(30%)	24	(30%)	13	(27%)	7	(9%)	30	(53%)***	20	(30%)	17	(25%)
Living alone n (%)	61	(46%)	30	(52%)	31	(41%)	41	(51%)	18	(37%)	32	(42%)	29	(51%)	33	(49%)	28	(42%)
Cluster A n (%)	56	(42%)	19	(33%)	37	(49%)	30	(37%)	24	(49%)	15	(19%)	41	(72%)***	24	(36%)	32	(48%)
Cluster B (other than BPD) n (%)	69	(51%)	27	(47%)	42	(55%)	43	(53%)	25	(51%)	29	(38%)	40	(70%)***	37	(55%)	32	(48%)
Cluster C n (%)	51	(38%)	23	(40%)	28	(37%)	33	(41%)	16	(38%)	15	(19%)	36	(63%)***	19	(28%)	32	(48%)*
Unemployed n (%)	95	(71%)	38	(66%)	57	(75%)	51	(63%)	42	(86%)**	53	(69%)	42	(74%)	48	(72%)	47	(70%)
Early loss n (%)	65	(49%)	28	(48%)	37	(49%)	41	(51%)	21	(43%)	38	(49%)	27	(47%)	31	(46%)	34	(51%)
Sexual abuse n (%)	69	(51%)	29	(50%)	40	(53%)	45	(56%)	24	(49%)	37	(48%)	32	(56%)	33	(49%)	36	(54%)
Physical abuse n (%)	52	(39%)	25	(43%)	27	(36%)	34	(42%)	17	(35%)	27	(35%)	25	(44%)	27	(40%)	25	(37%)
Alcohol use > 30 units/week n (%)	89	(66%)	35	(60%)	54	(71%)	52	(64%)	34	(69%)	47	(61%)	42	(74%)	45	(67%)	44	(66%)
Drug use > 4 times/week n (%)	91	(68%)	34	(59%)	57	(75%)	53	(65%)	34	(69%)	53	(69%)	38	(67%)	43	(64%)	48	(72%)

^{*}p < .05

^{**}p < .01

^{***}p < .001