



Data supplement to Redmore et al. Analysis of trends in adolescent suicides and accidental deaths in England and Wales, 1972-2011. Br J Psychiatry doi: 10.1192/bjp.bp.114.162347

Table DS1 International Classification of Diseases (ICD) coding ranges used to define specific methods of suicide and accidental death in England and Wales, 1972-2011.


Methods of suicide		
Method of suicide	ICD version^a	ICD codes
Solid or liquid poisoning	ICD-8 ICD-9 ICD-10	E950, E980 E950, E980 X60-65, X68-69, Y10-15, Y18-19
Gas poisoning	ICD-8 ICD-9 ICD-10	E951-952, E981-982 E951-952, E981-982 X66-67, Y16-17
Hanging, suffocation and strangulation	ICD-8 ICD-9 ICD-10	E953, E983 E953, E983 X70, Y20
Drowning	ICD-8 ICD-9 ICD-10	E954, E984 E954, E984 X71, Y21
Firearms and explosives	ICD-8 ICD-9 ICD-10	E955, E985 E955, E985 X72-75, Y22-25
Cutting and piercing instruments	ICD-8 ICD-9 ICD-10	E956, E986 E956, E986 X78, Y28
Jumping or falling from high places	ICD-8 ICD-9 ICD-10	E957, E987 E957, E987 X80, Y30
Other unspecified methods	ICD-8 ICD-9 ICD-10	E958-959, E988-989 E958-959, E988-989 X76-77, X79, X81-84, Y26-27, Y29, Y31-34
Methods of accidental death		
Poisoning	ICD-8 ICD-9 ICD-10	E850-877 E850-869 X40-49
Breathing; hanging, strangulation and suffocation	ICD-8 ICD-9 ICD-10	E913 E913 W75-77, W81, W83, W84

a. World Health Organisation International Statistical Classification of Diseases and Related Health Problems. ICD-8, 1972-1978; ICD-9, 1979-2000; ICD-10, 2001-2011.



Supplementary Figure DS1. Rates of suicide (with 95% confidence intervals) by solid and liquid poisoning, and hanging, in males and females aged (a) 10-14 years and (b) 15-19 years in England and Wales, 1972-2011

(a)



(b)




Figure DS2. Rates (per 100,000) of suicide (suicide and undetermined) deaths and accidental (poisoning and hanging) deaths for males and females aged: (a) 10-14 years and (b) 15-19 years in England and Wales, 1972 to 2011.