

Data supplement to Chisholm et al. Estimating the cost of implementing district mental healthcare plans in five low- and middle-income countries: the PRIME study. Br J Psychiatry doi: 10.1192/bjp.bp.114.153866

Table DS1 District-level contextualisation of core PRIME intervention packages

		Package localisation									
		Ethiopia		India		Nepal		South Africa		Uganda	
		HealthCare Visits	Drug Costs	HealthCare Visits	Drug Costs	HealthCare Visits	Drug Costs	HealthCare Visits	Drug Costs	HealthCare Visits	Drug Costs
Depression	Basic psychosocial intervention	X		N/C		N/C		X		X	
	Antidepressant medication	X	X	N/C	N/C	N/C	X	X	X	X	X
	Intensive psychosocial intervention					X		X			
	Psychosocial care for perinatal depression	N/C		N/C						X	
Psychosis	Basic psychosocial intervention	X		N/C		N/C		N/C		X	
	Antipsychotic medication	X	X	N/C		N/C	X	X	X	N/C	N/C
	Intensive psychosocial intervention					X		X			
Epilepsy	Basic psychosocial intervention	X				N/C				X	
	Anti-epileptic medication	X	X			N/C	X			X	N/C

Alcohol use Disorder	Identification and assessment	N/C		N/C		N/C		N/C		N/C	
	Brief interventions and follow-up	N/C		N/C		N/C		X		N/C	
	Management of alcohol withdrawal	X	X					X	X	N/C	N/C

Key	
X	Localisation change from tool default
N/C	No change from tool default
	Not Applicable
	Intervention not chosen by country