

Data supplement

Table DS1 Age-standardised rate of self-harm per 100 000 population and 95% confidence intervals, for persons aged 15+ years, in the three centres^a

	Oxford (n = 3751)		Manchester (n = 13 656) ^b		Derby (n = 6461)	
	Age-standardised rate	95% CI	Age-standardised rate	95% CI	Age-standardised rate	95% CI
Combined						
2000	338	309–369	508	473–544	432	401–466
2001	343	314–375	496	462–531	469	436–506
2002	383	353–415	464	431–500	482	448–518
2003	439	408–474	497	463–532	492	457–528
2004	359	330–391	432	401–466	439	408–474
2005	370	340–402	428	396–462	423	391–457
2006	328	300–359	419	388–453	413	382–446
2007	317	290–346	405	369–445	382	352–414
Average, 2000–2007	360	348–371	456	448–464	442	431–452
Males						
2000	290	254–330	426	382–477	364	325–408
2001	332	293–377	419	376–469	381	340–427
2002	339	299–384	356	317–402	432	387–483
2003	358	318–402	423	379–473	439	394–488
2004	286	249–326	335	296–380	357	317–402
2005	336	296–381	349	310–393	350	311–394
2006	271	235–312	334	294–379	357	317–402
2007	267	231–307	324	279–376	306	270–347
Average, 2000–2007	310	294–325	371	361–381	373	359–387
Females						
2000	387	345–434	589	537–647	499	452–551
2001	354	314–398	575	524–631	558	507–615
2002	431	386–482	574	523–631	531	482–587
2003	525	476–580	572	521–628	546	496–600
2004	437	393–485	533	583–589	522	473–576
2005	405	361–454	509	462–562	497	451–549
2006	387	345–434	510	462–563	471	423–524
2007	372	332–417	492	437–553	459	413–510
Average, 2000–2007	412	395–429	544	533–556	510	494–527

a. Age-standardised to the European population per 100 000. The 95% confidence intervals were calculated assuming a Poisson distribution. Local authority areas: Oxford City, City of Manchester, Derby Unitary Area.

b. Adjusted by 1.42 factor for period 2000 to end August 2002 to account for non-assessed episodes that were not collected.