

## **Data supplement**

Reference	Country	Period	n	Peak	Trough	Detail	Method
Watkins (2002) <sup>6</sup>	England	1971–1990 (born)	259	NS	NS	Excess in April–June births Great Ormond Street criteria <sup>16</sup>	Roger's chi-squared
Eagles (2001) <sup>4</sup>	Scotland	1965–1997 (recruited)	446	First half of year	Second half of year	Excess mainly in April–June Criteria not stated	Hewitt's test + chi-squared
Waller (2002) <sup>7</sup>	England		195	NS	NS	Only adult patients  Excess in April–June births in restrictive subtypes (n = 117)  DSM–IV <sup>17</sup> criteria	Chi-squared, Mann-Whitney and Spearman's test
Button (2007) <sup>5</sup>	England	After 1984 (recruited)	393	NS	NS	March peak not significant DSM–III <sup>18</sup> and DSM–IV criteria	Chi-squared + goodness of fit

## **Additional references**

- 16 Bryant-Waugh R, Lask B. Eating disorders in children. J Child Psychol Psychiatry 1995; 36: 191–202.
- 17 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (4th edn) (DSM-IV)*. APA, 1994.
- 18 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (3rd edn) (DSM–IIII)*. APA, 1980.