

Data supplement

Table DS1 Demographic, clinical, psychosocial functioning and quality of life characteristics according to threshold score on the Hypomania Checklist-32 and Bipolar Spectrum Diagnostic Scale: major depressive disorder with manic symptoms group (MDD-manic symptoms group) v. MDD group

	Hypomania Checklist-32			Bipolar Spectrum Diagnostic Scale		
	MDD-manic symptoms group (score 18 or more) (n = 42)	MDD group (score 17 or less) (n = 103)	P	MDD-manic symptoms group (score 12 or more) (n = 78)	MDD group (score 11 or less) (n = 67)	P
Age, years: mean (s.d.)	38.7 (9.5)	46.7 (12.8)	0.0004 ^a	41.7 (11.4)	47.6 (12.9)	0.004 ^a
Male, n (%)	19 (45)	27 (26)	0.03 ^b	29 (37)	17 (25)	0.13 ^b
Ethnicity, White: n (%)	41 (98)	100 (97)	0.86 ^b	76 (97)	65 (97)	0.88 ^b
Currently employed, n (%)	19 (45)	41 (40)	0.55 ^b	32 (41)	28 (42)	0.93 ^b
Premorbid IQ (NART), median (range)	35 (4–54)	31 (4–49)	0.14 ^c	32 (4–54)	34 (4–48)	0.58 ^c
Age at onset of depression, years: mean (s.d.)	20.6 (8.1)	26.6 (14.2)	0.01 ^a	22.7 (12.0)	27.3 (13.7)	0.03 ^a
MADRS, median (range)	17.5 (0–35)	10.0 (0–49)	0.01 ^c	17 (0–40)	7 (0–49)	0.00004 ^c
YMRS, median (range)	0 (0–5)	0 (0–5)	0.04 ^c	0 (0–5)	0 (0–5)	0.03 ^c
Chronic depression, n (%)	20 (48)	58 (56)	0.34 ^b	34 (44)	44 (66)	0.008 ^b
Episodes of depression, ^d median (range)	6 (3–40)	3 (1–30)	0.003 ^c	6 (1–40)	3 (1–30)	0.02 ^c
Ever admitted, n (%)	4 (10)	12 (12)	0.71 ^b	8 (10)	8 (12)	0.75 ^b
MGH score, median (range)	1.5 (0–8)	1.5 (0–8)	0.97 ^c	1.5 (0–10)	1.5 (0–6)	0.13 ^c
On psychotropic medication, n (%)	35 (83)	79 (77)	0.38 ^b	62 (80)	52 (78)	0.78 ^b
First-degree relative with depression, n (%)	23 (55)	47 (46)	0.32 ^b	40 (51)	30 (45)	0.44 ^b
First-degree relative with bipolar disorder, n (%)	4 (10)	3 (3)	0.09 ^b	3 (4)	4 (6)	0.55 ^b
Comorbid panic disorder, n (%)	18 (31)	25 (24)	0.41 ^b	27 (35)	11 (16)	0.01 ^b
Comorbid alcohol misuse, n (%)	23 (55)	18 (18)	0.00001 ^b	29 (37)	12 (18)	0.01 ^b
Comorbid alcohol dependence, n (%)	17 (41)	7 (7)	0.00001 ^b	19 (24)	5 (8)	0.006 ^b
GAF score, mean (s.d.)	63.4 (14)	69.0 (12.4)	0.001 ^a	64.1 (12.3)	71.2 (13.1)	0.001 ^a
FAST, median (range)						
Total score	31 (1–70)	24 (0–62)	0.02 ^c	31 (1–70)	18 (0–57)	0.0001 ^c
Autonomy	3 (0–10)	2 (0–12)	0.15 ^c	3 (0–11)	1 (0–12)	0.001 ^c
Occupational	6.5 (0–15)	4 (0–15)	0.03 ^c	6.5 (0–15)	3 (0–15)	0.002 ^c
Cognitive	7.5 (0–15)	6 (0–15)	0.10 ^c	8 (0–15)	5 (0–14)	0.009 ^c
Finances	2 (0–6)	0 (0–6)	0.02 ^c	2 (0–6)	0 (0–6)	0.002 ^c
Relationships	8 (0–18)	4 (0–18)	0.003 ^c	7 (0–18)	3 (0–18)	0.00006 ^c
Leisure time	3 (0–6)	3 (0–6)	0.56 ^c	3 (0–6)	3 (0–6)	0.09 ^c
WHOQOL-Bref, median (range)						
Total score	45.5 (19–69)	51 (29–73)	0.007 ^c	47 (19–68)	55 (29–73)	0.00002 ^c
Physical health	44 (6–81)	56 (6–94)	0.29 ^c	44 (6–81)	63 (6–94)	0.003 ^c
Psychological	34.5 (6–81)	50 (6–94)	0.0005 ^c	38 (6–81)	56 (6–94)	0.00001 ^c
Social relationships	44 (0–100)	56 (0–100)	0.02 ^c	50 (0–100)	56 (0–100)	0.002 ^c
Environment	56 (0–75)	63 (25–88)	0.007 ^c	56 (0–88)	69 (25–88)	0.00003 ^c

NART, National Adult Reading Test; MADRS, Montgomery-Åsberg Depression Rating Scale; YMRS, Young Mania Rating Scale; MGH, Massachusetts General Hospital;

a. Independent t-test.

b. χ^2 -test.

c. Mann-Whitney U-test.

d. For episodes of depression: sample size n = 78.