

## Data supplement

**Table DS1** Predictors of major depressive episode, post-traumatic stress disorder, generalised anxiety disorder and suicide risk among all participants

	Major depressive episode (n = 206)		Post-traumatic stress disorder (n = 206)		Generalised anxiety disorder (n = 206)		Suicide risk (n = 206)	
	Adjusted OR	(95% CI)	Adjusted OR	(95% CI)	Adjusted OR	(95% CI)	Adjusted OR	(95% CI)
Age, years	0.96	(0.91–1.01)	1.02	(0.96–1.07)	0.99	(0.94–1.04)	1.00	(0.95–1.05)
Years of education	0.88	(0.80–0.98)	1.04	(0.94–1.15)	1.02	(0.92–1.13)	0.97	(0.88–1.07)
Number of war-related traumatic events	1.01	(0.99–1.03)	1.01	(0.99–1.03)	1.01	(0.99–1.04)	1.03	(1.01–1.05)*
Torture (yes v. no)	1.13	(0.53–2.40)	0.74	(0.35–1.58)	0.75	(0.36–1.53)	1.26	(0.62–2.55)
Post-war death of relatives (yes v. no)	0.79	(0.26–2.37)	0.69	(0.18–2.69)	0.94	(0.35–2.55)	0.81	(0.31–2.15)
Monthly income	1.00	(1.00–1.00)	1.00	(1.00–1.00)	1.00	(1.00–1.00)	1.00	(1.00–1.00)
Group								
Widowed lone mothers	1 (Reference)		1 (Reference)		1 (Reference)		1 (Reference)	
Non-bereaved married mothers	0.08	(0.03–0.18)***	0.10	(0.04–0.23)***	0.14	(0.06–0.37)***	0.26	(0.12–0.61)**
Bereaved married mothers	0.12	(0.03–0.44)**	0.06	(0.01–0.28)***	0.34	(0.10–1.17)	0.34	(0.10–1.12)

OR, odds ratio.  
\* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ .