BJPsych
The British Journal of Psychiatry
1. doi: 10.1192/bjp.bp.111.099366

Data supplement

Table DS1 Observed and expected all-cause, cause-specific mortality and standardised mortality ratios (SMRs) among people who self-harm in Taipei City, January 2004 to December 2006

	Men ($n = 2316$)				Women ($n = 5285$)				Total ($n = 7601$)			
	First year		All follow-up years		First year		All follow-up years		First year		All follow-up years	
	O/E	SMR (95%CI)	O/E	SMR (95%CI)	O/E	SMR (95%CI)	O/E	SMR (95%CI)	O/E	SMR (95%CI)	O/E	SMR (95%CI)
All causes	172/13.9	12.4 (10.6–14.3)	273/39.9	6.8 (6.1–7.7)	113/11.2	10.1 (8.3–12.1)	243/34.9	7.0 (6.1–7.9)	285/25.2	11.3 (10.1–12.7)	516/74.8	6.9 (6.3–7.5)
Suicide	67/0.5	141.4 (109.5–179.5)	92/1.5	60.2 (48.5–73.8)	57/0.6	101.3 (76.7–131.2)	109/1.9	57.9 (47.5–69.8)	124/1.0	119.6 (99.5–142.6)	201/3.4	58.9 (51.0–67.6)
Non-suicide mortality	105/13.4	7.8 (6.4–9.5)	181/38.4	4.7 (4.1–5.5)	56/10.7	5.3 (4.0–6.8)	134/33.0	4.1 (3.4–4.8)	161/24.1	6.7 (5.7–7.8)	315/71.4	4.4 (3.9–4.9)
Malignant neoplasms	42/4.1	10.3 (7.4–13.9)	55/11.9	4.6(3.5–6.0)	13/3.6	3.6 (1.9–6.2)	26/11.3	2.3 (1.5–3.4)	55/7.6	7.2 (5.4–9.4)	81/23.2	3.5 (2.8–4.3)
Heart disease	7/1.7	4.0 (1.6–8.3)	17/4.9	3.5 (2.0–5.6)	5/1.1	4.5 (1.4–10.4)	16/3.4	4.7 (2.7–7.7)	12/2.8	4.2 (2.2–7.4)	33/8.3	4.0 (2.7–5.6)
Cerebrovascular disease	8/1.2	6.6 (2.8–13.0)	17/3.4	4.9 (2.9–7.9)	4/1.0	4.1 (1.1–10.6)	8/2.9	2.7 (1.2–5.4)	12/2.2	5.5 (2.8–9.6)	25/6.4	3.9 (2.5–5.8)
Diabetes mellitus	6/0.7	8.0 (2.9–17.5)	13/2.1	6.1 (3.2–10.4)	2/0.8	2.5 (0.3–9.0)	11/2.4	4.5 (2.3–8.1)	8/1.5	5.2 (2.2–10.2)	24/4.6	5.2 (3.4–7.8)
Pneumonia	2/0.7	2.8 (0.3–10.1)	9/1.9	4.7 (2.2–9.0)	1/0.4	2.7 (0.0–14.9)	1/1.1	0.9 (0.0–5.0)	3/1.1	2.8 (0.6–8.0)	10/3.0	3.3 (1.6–6.1)
Chronic liver disease and cirrhosis	0/0.3	0.0	3/0.9	3.3 (0.7–9.6)	1/0.2	5.0 (0.1–28.0)	3/0.6	4.8 (1.0–14.1)	1/0.5	2.0(0.0–11.2)	6/1.5	3.9 (1.4–8.5)
Nephritis, nephrotic syndrome and nephrosis	3/0.5	6.3 (1.3–18.5)	5/1.3	3.8 (1.2–8.9)	3/0.4	6.8 (1.4–20.0)	3/1.3	2.2 (0.5–6.6)	6/0.9	6.6 (2.4–14.3)	8/2.7	3.0 (1.3–5.9)
Hypertensive disease	0/0.1	0.0	1/0.4	2.6 (0.0–14.6)	1/0.1	7.7 (0.1–43.0)	3/0.4	7.8 (1.6–22.7)	1/0.3	3.7 (0.0–20.7)	4/0.8	5.2 (1.4–13.3)
Accident	8/0.5	14.7 (6.3–28.9)	14/1.8	8.0 (4.4–13.4)	3/0.3	9.5 (1.9–27.8)	8/1.0	7.7 (3.3–15.1)	11/0.9	12.8 (6.4–22.9)	22/2.8	7.9 (4.9–11.9)
Accidental poisoning	4/0.03	125.0 (33.6–320.0)	6/0.1	56.3 (20.6–122.6)	2/0.03	59.9 (6.7–216.3)	4/0.1	35.5 (9.6–90.9)	6/0.07	91.8 (33.5–199.8)	10/0.2	45.6 (21.8–83.9)
Other accident	4/0.5	7.8 (2.1–20.0)	8/1.6	4.9 (2.1–9.6)	1/0.3	3.6 (0.1–19.7)	4/0.9	4.3 (1.2–11.0)	5/0.8	6.3 (2.0–14.7)	12/2.6	4.7 (2.4–8.1)
Others	29/3.5	8.3 (5.5–11.9)	47/9.7	4.8 (3.5–6.4)	23/2.8	8.3 (5.3–12.5)	55/8.5	6.5 (4.9–8.5)	52/6.3	8.3 (6.2–10.9)	102/18.2	5.6 (4.6–6.8)

O, observed death; E, expected death.