

Data supplement

Table DS1 Symptoms stratified by low- and high-combat exposure^a

	Low-combat exposure						High-combat exposures					
	UK (<i>n</i> = 181)		US (<i>n</i> = 694)		UK v. US		UK (<i>n</i> = 132)		US (<i>n</i> = 866)		UK v. US	
	<i>n</i>	%	<i>n</i>	%	χ^2 (d.f.)	<i>P</i>	<i>n</i>	%	<i>n</i>	%	χ^2 (d.f.)	<i>P</i>
Physical symptoms												
Stomach pain	5	2.8	9	1.3	2.1 (1)	0.152	5	3.8	38	4.4	0.1 (1)	0.735
Headaches	21	11.9	50	7.2	4.0 (1)	0.045	20	15.2	120	13.9	0.1 (1)	0.709
Chest pain	9	5.1	17	2.5	3.4 (1)	0.067	9	6.8	46	5.3	0.5 (1)	0.490
Dizziness	3	1.7	9	1.3	0.2 (1)	0.688	4	3.1	12	1.4	2.0 (1)	0.159
Rapid heartbeat	5	2.8	11	1.6	1.2 (1)	0.274	5	3.8	51	5.9	1.0 (1)	0.319
Shortness of breath	5	2.8	21	3.0	0.02 (1)	0.881	6	4.6	48	5.6	0.2 (1)	0.625
Constipation/bowel problems	3	1.7	26	3.8	2.0 (1)	0.161	5	3.8	68	6.8	1.7 (1)	0.194
Nausea/indigestion	16	8.8	40	5.8	2.2 (1)	0.137	5	3.8	73	8.5	3.5 (1)	0.062
Trouble sleeping	37	20.9	117	17.0	1.4 (1)	0.230	31	23.4	278	32.3	4.1 (1)	0.043
Physical symptoms binary 2 + physical symptoms	23	12.7	72	10.4	0.8 (1)	0.369	22	16.7	187	21.6	1.7 (1)	0.195

a. Unadjusted for demographic differences between the two samples. Numbers may not add up to total because of missing values.