

Potential host range of *Cotesia vanessae* (Hymenoptera: Braconidae), a parasitoid new to North America and a possible biological control agent of noctuid pest species — Supplementary Table 1

V.A.D. Hervet, R.A. Laird, and K.D. Floate

Supplementary Table 1. Fate of non-host Lepidoptera in which *C. vanessae* oviposited, but from which no F₁ progeny emerged.

Lepidoptera species	Number of caterpillars that:	
	Survived	Died
FAMILY DREPANIDAE		
<i>Habrosyne scripta</i> (Gosse)	27	3
FAMILY EREBIDAE		
<i>Anticarsia gemmatalis</i> Hübner	39	6
<i>Spilosoma virginica</i> (Fabricius)	30	0
FAMILY NOCTUIDAE		
<i>Lacinipolia renigera</i> (Stephens)	22	3
<i>Lacinipolia sareta</i> (Smith)	31	3
<i>Spodoptera eridania</i> (Stoll)	19	3
<i>Spodoptera exigua</i> (Hübner)	27	2
<i>Spodoptera frugiperda</i> (J. E. Smith)	5	13
<i>Trichordestra lilacina</i> (Harvey)	19	39
<i>Xestia c-nigrum</i> (Linnaeus)	17	16
<i>Sideridis rosea</i> (Harvey)	26	9
FAMILY PIERIDAE		
<i>Pieris rapae</i> (Linnaeus)	12	27
FAMILY SPHINGIDAE		
<i>Hyles euphorbiae</i> (Linnaeus)	12	5