

Geological Magazine											
The Significance of Iron-Stained Foraminifera off SE Trinidad, West Indies, Western Central Atlantic Ocean											
Brent Wilson											
Appendix 1: Unstained calcareous benthonic foraminifera from the Savonette Field.											
Species (specimens not iron-stained)	Station 1	Station 2	Station 3	Station 4	Station 5	Station 6	Station 7	Station 8	Station 9	Station 10	Station 11
<i>Ammonia pauciloculata</i>	0	0	0	0	1	0	0	0	0	0	0
<i>Ammonia tepida</i>	0	0	0	0	0	0	0	0	0	0	1
<i>Amphicoryna sublineata</i>	1	0	3	0	1	0	1	0	1	0	1
<i>Amphistegina gibbosa</i>	10	7	1	6	1	2	1	2	0	2	37
<i>Amphistegina radiata</i>	0	1	0	0	0	0	0	0	0	0	1
<i>Angulogerina occidentalis</i>	6	6	0	0	0	0	0	0	0	0	0
<i>Anomalinoidea sp.</i>	1	0	0	0	0	0	0	0	0	0	0
<i>Brizalina barbata</i>	0	0	0	2	1	0	0	0	0	0	1
<i>Brizalina fragilis</i>	0	0	0	0	0	0	0	1	0	0	1
<i>Brizalina striatula spinata</i>	1	1	0	0	0	0	0	1	0	1	0
<i>Buccella hannai</i>	1	0	0	0	0	0	0	0	0	0	0
<i>Bulimina striata mexicana</i>	0	0	0	0	0	0	0	0	1	0	0
<i>Cancris sagra</i>	0	0	1	0	2	0	0	2	0	1	0
<i>Caribbeanella polystoma</i>	0	0	0	0	1	1	0	0	0	0	0
<i>Cassidulina laevigata</i>	0	0	1	0	0	0	0	0	0	0	0
<i>Cassidulina norcrossi australis</i>	5	3	7	5	11	6	4	5	3	2	0
<i>Cibicides ex gr. pseudoungerianus</i>	42	43	40	42	52	26	40	59	39	56	59
<i>Cibicides floridanus</i>	3	0	0	4	3	2	1	0	0	0	0
<i>Cibicides io</i>	10	22	35	20	33	18	19	23	23	14	4
<i>Cibicides sp. 1</i>	0	0	0	0	1	2	0	0	1	0	0
<i>Cibicides sp. A Poag 1981</i>	0	0	0	0	0	0	1	0	0	0	0
<i>Dentalina communis</i>	0	1	0	0	0	0	0	0	0	0	0
<i>Discorbis floridensis</i>	5	2	1	6	2	1	6	2	0	2	2
<i>Elphidium sp.</i>	0	1	1	0	0	0	0	0	0	0	0
<i>Elphidium translucens</i>	5	3	10	3	5	7	3	1	2	2	2
<i>Eponides antillarum</i>	15	20	11	20	6	10	8	0	6	10	15
<i>Eponides campester</i>	2	1	0	0	0	0	0	1	3	0	0
<i>Eponides regularis</i>	0	0	0	0	3	0	0	1	0	0	0
<i>Eponides repandus</i>	2	4	0	0	0	0	0	8	0	0	1
<i>Gavelinopsis sp.</i>	0	0	2	0	3	0	1	0	0	0	0
<i>Globocassidulina subglobosa</i>	3	7	5	3	9	8	2	9	6	7	2
<i>Gyroidina sp.</i>	0	1	1	0	0	0	0	0	0	0	0
<i>Hanzawaia concentrica</i>	8	9	13	6	5	7	0	11	4	3	4
<i>Heterostegina antillarum</i>	0	0	0	0	0	0	0	0	0	0	4
<i>Lenticulina atlantica</i>	0	0	0	0	0	0	0	0	0	1	0
<i>Lenticulina gibba</i>	0	0	0	0	0	0	0	0	2	0	0
<i>Lenticulina iota</i>	0	0	0	0	0	0	1	0	0	0	0
<i>Lenticulina occidentalis</i>	0	0	0	0	0	0	0	0	0	0	3
<i>Lenticulina rotulata</i>	2	2	1	0	1	1	1	3	0	2	0
<i>Marginulinopsis planatus</i>	2	6	2	3	1	3	5	0	1	1	17
<i>Miliolinella subrotunda</i>	0	2	2	0	1	0	0	1	0	0	2
<i>Neoconorbina sp.</i>	0	0	1	0	0	0	0	0	0	0	0
<i>Nodosaria pyrula</i>	0	0	0	0	0	1	0	0	0	0	0
<i>Nodosaria vertebralis</i>	0	0	0	0	1	0	0	1	0	0	0
<i>Pavonina atlantica</i>	0	0	0	0	0	0	0	0	0	0	1
<i>Planularia sp.</i>	0	0	0	1	0	0	0	0	0	0	0
<i>Pullenia quinqueloba</i>	0	1	1	0	0	0	0	1	0	0	0
<i>Quinqueloculina auberiana</i>	1	0	0	0	0	1	0	0	0	0	0
<i>Quinqueloculina bicostata</i>	2	1	0	0	1	0	0	0	0	0	2
<i>Quinqueloculina compta</i>	1	0	1	0	0	0	0	0	0	0	0
<i>Quinqueloculina lamarckiana</i>	0	2	1	2	1	2	1	0	0	0	6
<i>Quinqueloculina seminula</i>	0	0	0	1	0	0	1	0	0	0	0
<i>Quinqueloculina sp.</i>	0	0	0	0	0	0	0	1	0	0	0
<i>Reussella atlantica</i>	0	4	4	1	4	0	1	2	0	0	2
<i>Rosalina floridana</i>	0	0	0	0	0	1	0	0	0	0	0
<i>Rotalid indet.</i>	0	1	0	0	0	0	0	0	0	0	0
<i>Saracenaria arcuata</i>	0	0	0	0	1	0	0	0	0	0	0
<i>Siphonina pulchra</i>	1	0	2	2	3	4	1	0	3	0	2
<i>Stomatrobina concentrica</i>	0	0	1	0	2	0	0	0	0	0	0
<i>Trifarina bradyi</i>	1	2	2	0	1	1	2	0	0	1	1
<i>Triloculina tricarinata</i>	1	0	0	0	0	0	0	0	0	0	1
<i>Triloculina trigonula</i>	0	0	0	0	0	0	0	0	1	0	0
<i>Uvigerina auberiana</i>	0	0	0	0	0	0	0	0	0	1	1
<i>Uvigerina laevis</i>	0	2	1	0	1	0	0	0	0	0	0
<i>Uvigerina subperegrina</i>	0	0	1	0	0	0	0	0	0	0	0
Northings	1110575	1110820	1111482	1112739	1112745	1112722	1111511	1111477	1113728	1116206	1100352
Eastings	0801432	0800955	0800272	0799492	0800566	0801420	0801427	0800405	0799255	0797440	0799865
Water depth (metres)	87	87	87	87	87	90	89	87	87	85	78
Residue mass (grams)	7.48	5.53	7.29	11.01	3.46	6.34	8.55	3.04	2.11	3.99	4.78
Percentage of fauna as planktonics (%p)	57.50	25.00	31.00	26.50	27.00	38.50	35.50	39.00	26.50	18.50	25.00
Percentage of iron-stained planktonics	59.1	84.0	67.7	71.7	74.0	79.2	74.6	78.2	75.5	78.4	82.0
Number of fresh benthonic specimens	131	155	152	127	158	104	100	135	96	106	173
Information Function (H) for fresh benthonic assemblages	2.52	2.56	2.41	2.19	2.43	2.44	2.10	2.00	1.91	1.75	2.24