




Supplementary Figure 2: Case 1 – 24-year-old right hand dominant female patient with a history of 2 generalized seizures. MRI FLAIR sequence is shown. There was NO enhancement with gadolinium.



Supplementary Figure 3: Case 2 - 52-year-old right hand dominant male with focal aware seizures with motorized aphasia. Neurologically intact. MRI FLAIR sequence is shown. There was NO enhancement with gadolinium.



Supplementary Figure 4: Case 3 - 49-year-old right hand dominant male with focal unaware seizures without any neurological deficit. MRI FLAIR sequence is shown. There was NO enhancement with gadolinium.



Please be advised that the supplementary figures were case presentations obtained from the following article:

Khan OH, Mason W, Kongkham PN, Bernstein M, Zadeh G. Neurosurgical management of adult diffuse low grade gliomas in Canada: a multi-center survey. *J Neurooncol*. 2016;126:137-149.
doi:10.1007/s11060-015-1949-0

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.