

Variable	Discharge Functional Status (probability of dependence)		Discharge Destination (ordinal probability of worse outcome)		Length of Admission (computed mean estimate)	
	Estimate (95% CI)	p-value	Estimate (95% CI)	p-value	Mean Estimate (Std Error)	p-value
Age (per 10 year increment)	1.30 (1.09-1.56)	0.0046	1.24 (1.08-1.43)	0.0025	11.48% increase in LOS	0.0002
History of Stroke or TIA	---	---	1.23 (0.79-1.91)	0.3578	---	---
Baseline Functional Dependence	---	---	2.07 (1.22-3.52)	0.0075	---	---
Onset on Waking (Yes)	---	---	1.31 (0.78-2.21)	0.3036	27.24 (3.09)	0.0028
Onset on Waking (No)	---	---	---	---	19.80 (1.65)	
Stroke Severity (Severe)	---	---	---	---	25.75 (2.17)	0.0122
Stroke Severity (Moderate)	---	---	---	---	20.95 (2.14)	
Presentation < 3 hours	---	---	---	---	27.49 (1.96)	0.0279
Presentation 3 - < 6 hrs	1.71 (0.72-4.03)	0.36	1.29 (0.72-2.34)	0.3942	26.80 (3.23)	
Presentation 6 - < 12 hrs	4.63 (0.60-35.68)	0.4546	0.43 (0.13-1.40)	0.1615	15.15 (2.75)	
Presentation > 12 hrs	5.31 (1.23-23.02)	0.2293	1.14 (0.65-2.00)	0.6444	26.05 (3.31)	
No Treatment	---	---	---	---	36.08 (2.48)	< 0.001
Use of tPA Alone	0.48 (0.26-0.89)	0.0201	0.48 (0.31-0.75)	0.0013	27.23 (2.78)	
Use of EVT Alone	0.36 (0.12-1.08)	0.0679	0.18 (0.07-0.46)	0.0004	14.00 (2.60)	
Use of tPA and EVT	0.31 (0.15-0.66)	0.0023	0.32 (0.17-0.59)	0.0003	21.14 (2.81)	

Supplemental Table 1 - Multivariate analysis of variables associated with secondary outcomes: discharge functional status, discharge destination, and length of admission.