

Lithostratigraphy, age and distribution of Eocene volcanic sequences on eastern King George Island, South Shetland Islands, Antarctica

Smellie, J.L.^{1*}, Hunt, R.J.², McIntosh, W.C.³ and Esser, R.P.^{3,4}

¹ School of Geography, Geology & the Environment, University of Leicester, Leicester LE1 7RH, UK [jls55@le.ac.uk]

² School of Earth Sciences, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT [rjh138@icloud.com]

³ New Mexico Bureau of Geology and Mineral Resources, New Mexico Tech, Socorro, NM, 87801, USA [mcintosh@nmt.edu]

⁴ now at: Energy and Geosciences Institute, University of Utah, Salt Lake City, UT, 84108, USA [richard.esser@utah.edu]

Supplementary Information file S1 – exposure maps for each of the areas studied on King George Island




Fig. S1-1. Point Hennequin—Warkocz area




Fig. S1-2. Rembiszewski Nunataks area




Fig. S1-3. Vauréal Peak—Puchalski Peak area




Fig. S1-4. Harnasie Hill area




Fig. S1-5. Point Thomas—Sphinx Hill area




Fig. S1-6. Godwin Cliffs—Lions Rump—Low Head area