

A review of preservational variation of fossil inclusions in amber of different chemical groups

Victoria E. McCoy, Carmen Soriano & Sarah E. Gabbott

References for Supplementary Tables 1 & 2

- Akhmetiev, M. A. 2010. Paleocene and Eocene floristic and climatic change in Russia and Northern Kazakhstan. *Bulletin of Geosciences* **85**, 17-34.
- Andersen, N. M. & Poinar, G. 1992. Phylogeny and classification of an extinct water strider genus (Hemiptera, Gerridae) from Dominican amber, with evidence of mate guarding in a fossil insect. *Journal of Zoological Systematics and Evolutionary Research* **30**, 256-267.
- Archibald, S. B. 2005. New Dinopanorpidae (Insecta: Mecoptera) from the Eocene Okanagan Highlands (British Columbia, Canada and Washington State, USA). *Canadian Journal of Earth Sciences* **42**, 119-136.
- Azar, D., Adaymeh, C. & Jreich, N. 2007. Paleopsychoda zherikhini, a new Cretaceous species of moth flies from Taimyr amber (Diptera: Psychodidae: Psychodinae). *African Invertebrates* **48**, 163-168.
- Azar, D., Gèze, R. & Acra, F. 2010. Lebanese amber. In Penney, D. (ed.) *Biodiversity of fossils in amber from the major world deposits*, 271-298. Manchester, UK: Siri Scientific Press.
- Bauer, A.M., Böhme, W. & Weitschat, W. 2005. An Early Eocene gecko from Baltic amber and its implications for the evolution of gecko adhesion. *Journal of Zoology* **265**, 327-332.
- Blagoderov, V. & Grimaldi, D. 2004. Fossil Sciaroidea (Diptera) in Cretaceous ambers, exclusive of Cecidomyiidae, Sciaridae, and Keroplatidae. *American Museum Novitates* 3433, 1-76.
- Bosselaers, J. 2004. A new *Garcorops* species from Madagascar copal (Araneae: Selenopidae). *Zootaxa* **445**, 1-7.
- Brandão, C. R. F., Urbani, C. B., Wagensberg, J. & Yamamoto, C. I. 1998. New *Technomyrmex* in Dominican amber (Hymenoptera: Formicidae), with a reappraisal of Dolichoderinae phylogeny. *Insect Systematics & Evolution* **29**, 411-428.
- Brasier, M., Cotton, L. & Yenney, I. 2009. First report of amber with spider webs and microbial inclusions from the earliest Cretaceous (c.140 Ma) of Hastings, Sussex. *Journal of the Geological Society* **166**, 989-997.
- Breton, G. 2010 Les Actinomycetales de l'ambre Sparnacien des Corbières (Aude, France): taphonomie et diversité. *Annales de la Société géologique du Nord* **17**, 3-22.
- Butcher, B. A., Zaldivar-Riveron, A., Van De Kamp, T., Dos Santos Rolo, T., Baumbach, T. & Quicke, D. L. 2014. Extension of historical range of Betylobraconinae (Hymenoptera: Braconidae) into Palaeartic Region based on a Baltic amber fossil, and description of a new species of *Mesocentrus Szépligeti* from Papua New Guinea. *Zootaxa* **3860**, 449-463.
- Calvillo-Canadell, L., Cevallos-Ferriz, S. R. & Rico-Arce, L. 2010. Miocene Hymenaea flowers preserved in amber from Simojovel de Allende, Chiapas, Mexico. *Review of Palaeobotany and Palynology* **160**, 126-134.
- Chatzimanolis, S., Newton, A. F., Soriano, C., & Engel, M. S. 2013. Remarkable stasis in a phloeocharine rove beetle from the Late Cretaceous of New Jersey (Coleoptera, Staphylinidae): *Journal of Paleontology* **87**, 177-182.
- Choufani, J., Azar, D., Perrichot, V., Soriano, C., Tafforeau, P. & Nel, A. 2011. The genus *Leptoconops* Skuse (Diptera: Ceratopogonidae) in Early Cretaceous Charentese amber. *Palaeobiodiversity and Palaeoenvironments* **91**, 285-291.
- Choufani, J., Perrichot, V., Azar, D. & Nel, A. 2014. New biting midges (Diptera: Ceratopogonidae) in late Cretaceous Vendean amber. *Paleontological Contributions* **10H**, 34-40.

- Choufani, J., El-Halabi, W., Azar, D. & Nel, A. 2015. First fossil insect from Lower Cretaceous Lebanese amber in Syria (Diptera: Ceratopogonidae). *Cretaceous Research* **54**, 106-116.
- Conran, J., Busch, A., Engel, M. S., Kaulfuss, U., Kiecksee, A. P., Lee, D., Maraun, M. & Perrichot, V. 2013. The first arthropods from New Zealand amber. *In Australian Entomological Society 44th Annual Meeting & Scientific Conference 2013-Invertebrates in extreme environments*, 64.
- Coty, D., Aria, C., Garrouste, R., Wils, P., Legendre, F. & Nel, A. 2014. The first ant-termite syninclusion in amber with CT-scan analysis of taphonomy. *PloS one* **9**, e104410.
- Dawson, M. R., West, R. M., Ramaekers, P. & Hutchison, J. H. 1975. New evidence on the palaeobiology of the Eureka Sound Formation, Arctic Canada. *Arctic* **28**, 110-116.
- Delclòs, X., Arillo, A., Peñalver, E., Barrón, E., Soriano, C., Del Valle, R. L., Bernárdez, E., Corral, C. & Ortuño, V. M. 2007. Fossiliferous amber deposits from the Cretaceous (Albian) of Spain. *Comptes Rendus Palevol* **6**, 135-149.
- Denton Jr, R. K. & O'Neill, R. C. 1995. *Prototeius stageri*, gen. et sp. nov., a new teiid lizard from the Upper Cretaceous Marshalltown Formation of New Jersey, with a preliminary phylogenetic revision of the Teiidae. *Journal of Vertebrate Paleontology* **15**, 235-253.
- Depalma, R., Cichocki, F., Dierick, M., & Feeney, R. 2010. Preliminary notes on the first recorded amber insects from the Hell Creek Formation. *The Journal of Paleontological Sciences* **10**, 1-7.
- Dierick, M., Cnudde, V., Masschaele, B., Vlassenbroeck, J., Van Hoorebeke, L. & Jacobs, P. 2007. Micro-CT of fossils preserved in amber. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **580**, 641-643.
- Dunlop, J. A., Penney, D., Dalüge, N., Jäger, P., Mcneil, A., Bradley, R. S., Withers, P. J., & Preziosi, R. F. 2011. Computed tomography recovers data from historical amber: an example from huntsman spiders. *Naturwissenschaften* **98**, 519-527.
- Dunlop, J. A., Wirth, S., Penney, D., McNeil, A., Bradley, R. S., Withers, P. J. & Preziosi, R. F. 2012. A minute fossil phoretic mite recovered by phase-contrast X-ray computed tomography. *Biology letters* **8**, 457-460.
- Dunlop, J. A., Selden, P. A., & Giribet, G. 2016. Penis morphology in a Burmese amber harvestman. *The Science of Nature* **103**, 1-5.
- Durán-Ruiz, C., Riquelme, F., Coutiño-José, M., Carbot-Chanona, G., Castaño-Meneses, G., Ramos-Arias, M. & Jin, J. 2013. Ants from the Miocene Totolapa amber (Chiapas, Mexico), with the first record of the genus *Forelius* (Hymenoptera, Formicidae). *Canadian Journal of Earth Sciences* **50**, 495-502.
- Edgecombe, G. D., Vahtera, V., Stock, S. R., Kallonen, A., Xiao, X., Rack, A. & Giribet, G. 2012. A scolopocryptopid centipede (Chilopoda: Scolopendromorpha) from Mexican amber: synchrotron microtomography and phylogenetic placement using a combined morphological and molecular data set. *Zoological Journal of the Linnean Society* **166**, 768-786.
- Engel, M. S., Nel, A., Azar, D., Soriano, C., Tafforeau, P., Neraudeau, D., Colin, J-P. & Perrichot, V. 2011. New, primitive termites (Isoptera) from Early Cretaceous ambers of France and Lebanon. *Palaeodiversity* **4**, 39-49.
- Engel, M. S., Ortega-Blanco, J., Soriano, C., Grimaldi, D. A. & Delclòs, X. 2013. A new lineage of enigmatic diaprioid wasps in Cretaceous amber (Hymenoptera: Diaprioidea). *American Museum Novitates* **3771**, 1-23.
- Fursov, V., Shirota, Y., Nomiya, T. & Yamagishi, K. 2002. New Fossil Mymarommatid Species, *Palaeomymar japonicum* sp. nov. (Hymenoptera: Mymarommatidae), Discovered in Cretaceous Amber from Japan (Systematics, Morphology and Evolution). *Entomological science* **5**, 51-54.
- Golovneva, L. & Shchepetov, S. 2010. Phytostratigraphy of Albian-Cenomanian sediments in the Kiya River basin (the Chulym-Yenisei area of the west Siberian lowland). *Stratigraphy and Geological Correlation* **18**, 153-165.

- Gomez, B., Bamford, M., Martínez-Delclòs, X. 2002. Lower Cretaceous plant cuticles and amber (Kirkwood Formation, South Africa). *Comptes Rendus Palevol* **1**, 83-87.
- Greco, M. K., Welz, P. M., Siegrist, M., Ferguson, S. J., Gallmann, P., Roubik, D. W. & Engel, M. S. 2011. Description of an ancient social bee trapped in amber using diagnostic radioentomology. *Insectes Sociaux* **58**, 487-494.
- Grimaldi, D., Bonwich, E., Delannoy, M. & Doberstein, S. 1994. Electron microscopic studies of mummified tissues in amber fossils. *American Museum Novitates* **3097**, 1-31.
- Grimaldi, D. & Case, G. R. 1995. A feather in amber from the Upper Cretaceous of New Jersey. *American Museum Novitates* **3126**, 1-6.
- Grimaldi, D. A. & Nascimbene, P. C. 2010. Raritan (New Jersey) amber. In Penney, D. (ed.) *Biodiversity of fossils in amber from the major world deposits*, 167-191. Manchester, UK: Siri Scientific Press.
- Hand, S., Archer, M., Bickel, D., Creaser, P., Dettmann, M., Godthelp, H., Jones, A., Norris, B. & Wicks, D. 2010. Australian cape york amber. In Penney, D. (ed.) *Biodiversity of fossils in amber from the major world deposits*, 69-79. Manchester, UK: Siri Scientific Press.
- Heethoff, M., Helfen, L. & Norton, R. A. 2009. Description of *Neoliodes dominicus* n. sp. (Acari, Oribatida) from Dominican Amber, aided by synchrotron X-ray microtomography. *Journal of Paleontology* **83**, 153-159.
- Henderickx, H., Cnudde, V., Masschaele, B., Dierick, M., Vlassenbroeck, J., & Van Hoorebeke, L. 2006. Description of a new fossil *Pseudogarypus* (Pseudoscorpiones: Pseudogarypidae) with the use of X-ray micro-CT to penetrate opaque amber. *Zootaxa* **1305**, 41-50.
- Henderickx, H., Tafforeau, P., & Soriano, C. 2012. Phase contrast synchrotron microtomography reveals the morphology of a partially visible new *Pseudogarypus* in Baltic amber (Pseudoscorpiones: Pseudogarypidae): *Palaeontologia Electronica* **15**, 11.
- Henderickx, H., Bosselaers, J., Pauwels, E., Van Hoorebeke, L. & Boone, M. 2013a. X-ray micro-CT reconstruction reveals eight antennomeres in a new fossil taxon that constitutes a sister clade to Dundoxenos and Triozocera (Strepsiptera: Corioxenidae). *Palaeontologia Electronica* **16**, 16-31.
- Henderickx, H., Perkovsky, E. E., Van Hoorebeke, L. & Boone, M. 2013b. The first pseudogarypid in Rovno amber (Ukraine) (Pseudoscorpiones: Pseudogarypidae). *PHEGGA* **41**, 90-92.
- Hinojosa-Diaz, I. A. & Engel, M. S. 2007. A new fossil orchid bee in Colombian copal (Hymenoptera: Apidae). *American Museum Novitates* **3589**, 1-7.
- Kasiński, J. & Tołkanowicz, E. 1999. Amber in the northern Lublin region. In: Kosmowska-Ceranowicz, B. & Paner, H., eds, *Investigation into Amber. Proceedings of the International Interdisciplinary Symposium: Baltic Amber and other Fossil Resins 997*. Urbs Gyddanyzc-1997. Gdańsk, pp. 41-51.
- Kedves, M. & Párdutz, Á. 2002. Ultrastructure of 100 million years old microorganisms of the Ajkaite (Upper Cretaceous amber) from Hungary. *TAIWANIA-TAIPEI* **47**, 273-280.
- Kehlmaier, C., Dierick, M., & Skevington, J. H. 2014. Micro-CT studies of amber inclusions reveal internal genitalic features of big-headed flies, enabling a systematic placement of *Metanephrocerus* Aczél, 1948 (Insecta: Diptera: Pipunculidae). *Arthropod Systematics and Phylogeny* **72**, 23-36.
- Kielhorn, K-H. & Blick, T. 2007. Erstfund von *Hahnia picta* (Araneae, Hahniidae) in Deutschland—mit Angaben zu Habitatpräferenz und Verbreitung. *Arachnologische Mitteilungen* **33**, 7-10.
- Kirejtshuk, A. G., Azar, D., Tafforeau, P., Boistel, R. & Fernandez, V. 2009. New beetles of Polyphaga (Coleoptera, Polyphaga) from Lower Cretaceous Lebanese amber. *Denisia* **26**, 119-130.
- Kononova, S. 2003. New species of egg-parasitizing wasps of the genus *Idris* (Scelionidae, Proctotrupeoidea) of the Rovno Amber. *Paleontological Journal c/c of Paleontologicheskii zhurna* **37**, 275-279.

- Koteja, J. 2004. Scale insects (Hemiptera: Coccinea) from Cretaceous Myanmar (Burmese) amber. *Journal of Systematic Palaeontology* **2**, 109-114.
- Lak, M., Azar, D., Nel, A., Neraudeau, D. & Tafforeau, P. 2008. The oldest representative of the Trichomyiinae (Diptera: Psychodidae) from the Lower Cenomanian French amber studied with phase-contrast synchrotron X-ray imaging. *Invertebrate Systematics* **22**, 471-478.
- Lak, M. & Nel, A. 2009. An enigmatic diapriid wasp (Insecta, Hymenoptera) from French Cretaceous amber. *Geodiversitas* **31**, 137-144.
- Lambert, J. B., Santiago-Blay, J. A. & Anderson, K. B. 2008. Chemical signatures of fossilized resins and recent plant exudates. *Angewandte Chemie International Edition* **47**, 9608-9616.
- Lambert, J., Tsai C. H., Shah, M., Hurlley, A. & Santiago-Blay, J. 2012. Distinguishing Amber and Copal Classes by Proton Magnetic Resonance Spectroscopy. *Archaeometry* **54**, 332-348.
- Lambert, J. B., Santiago-Blay, J. A., Wu, Y. & Levy, A. J. 2015. Examination of amber and related materials by NMR spectroscopy. *Magnetic Resonance in Chemistry* **53**, 2-8.
- Langenheim, R., Smiley, C. & Gray, J. 1960. Cretaceous amber from the Arctic coastal plain of Alaska. *Geological Society of America Bulletin* **71**, 1345-1356.
- Lazell Jr, J. D. 1965. An *Anolis* (Sauria, Iguanidae) in amber. *Journal of Paleontology* **39**, 379-382.
- Longrich, N. R. 2011. *Titanoceratops ouranos*, a giant horned dinosaur from the late Campanian of New Mexico. *Cretaceous Research* **32**, 264-276.
- Lyubarsky, G. Y. & Perkovsky, E. 2014. New species of the genus *Nganasania* from Upper Cretaceous of Taymyr (Coleoptera: Cryptophagidae). *Russian Entomological Journal* **23**, 191-194.
- Makarkin, V. N. & Kupryjanowicz, J. 2010. A New Mantispid-Like Species of Rhachiberothinae (Neuroptera: Berothidae) from Baltic Amber, with a Critical Review of the Fossil Record of the Subfamily. *Acta Geologica Sinica (English Edition)* **84**, 655-664.
- Martill, D. M., Loveridge, R. F., Andrade, D., Gomes, J. F. & Cardoso, A. H. 2005. An unusual occurrence of amber in laminated limestones: the Crato Formation Lagerstätte (Early Cretaceous) of Brazil. *Palaeontology* **48**, 1399-1408.
- Martínez-Delclòs, X., Briggs, D. E. & Peñalver, E. 2004. Taphonomy of insects in carbonates and amber. *Palaeogeography, Palaeoclimatology, Palaeoecology* **203**, 19-64.
- McAlpine, J. & Martin, J. 1969. Canadian amber—a paleontological treasure-chest. *The Canadian Entomologist* **101**, 819-838.
- McKellar, R. C. & Wolfe, A. P. 2010. Canadian amber. In Penney, D. (ed.) *Biodiversity of fossils in amber from the major world deposits*, 96-113. Manchester, UK: Siri Scientific Press.
- Michener, C. D. & Grimaldi, D. A. 1988. A *Trigona* from Late Cretaceous amber of New Jersey (Hymenoptera: Apidae: Meliponinae). *American Museum Novitates* **2917**, 1-10.
- Mustoe, G. E. 1985. Eocene amber from the Pacific coast of North America. *Geological Society of America Bulletin* **96**, 1530-1536.
- Najarro, M., Peñalver, E., Rosales, I., Pérez-De-La-Fuente, R., Daviero-Gomez, V., Gomez, B. & Delclòs, X. 2009. Unusual concentration of Early Albian arthropod-bearing amber in the Basque-Cantabrian Basi (El Soplao, Cantabria, Northern Spain): Palaeoenvironmental and palaeobiological implications. *Geologica Acta* **7**, 363-388.
- Nadein, K. S., Perkovsky, E. E., & Moseyko, A. G. 2016. New Late Eocene Chrysomelidae (Insecta: Coleoptera) from Baltic, Rovno and Danish ambers. *Papers in Palaeontology* **2**, 117-137.
- Nel, A., Depalma, R. A. & Engel, M. S. 2010. A possible hemiphlebiid damselfly in Late Cretaceous amber from South Dakota (Odonata: Zygoptera). *Transactions of the Kansas Academy of Science* **113**, 231-234.
- Nel, A. & Prokop, J. 2005. New Fossil Scelionidae (Insecta: Hymenoptera) in Early Paleogene Amber from Eastern Moravia (Czech Republic) and Northern France. *Polskie Pismo Entomologiczne* **74**, 339-347.

- Onoratini, G., Guiliano, M., Mille, G. & Simon, P. 2009. L'ambre albo-cénomanién de la montagne de Lure (Alpes-de-Haute-Provence), outil stratigraphique et paléogéographique. *Geobios* **42**, 89-99.
- Peñalver, E., Álvarez-Fernández, E., Arias, P., Delclòs, X. & Ontañón, R. 2007. Local amber in a Palaeolithic context in Cantabrian Spain: the case of La Garma A. *Journal of Archaeological Science* **34**, 843-849.
- Peñalver, E. & Delclòs, X. 2010. Spanish amber. In Penney, D. (ed.) *Biodiversity of fossils in amber from the major world deposits*, 236-270. Manchester, UK: Siri Scientific Press.
- Penney, D. 2010. Dominican amber. In Penney, D. (ed.) *Biodiversity of fossils in amber from the major world deposits*, 22-41. Manchester, UK: Siri Scientific Press.
- Penney, D., Dierick, M., Cnudde, V., Masschaele, B., Vlassenbroeck, J., Van Hoorebeke, L. & Jacobs, P. 2007. First fossil Micropholcommatidae (Araneae), imaged in Eocene Paris amber using X-ray computed tomography. *Zootaxa* **1623**, 47-53.
- Penney, D., Mcneil, A., Green, D. I., Bradley, R., Marusik, Y. M., Withers, P. J., & Preziosi, R.F. 2011. A new species of anapid spider (Araneae: Araneoidea, Anapidae) in Eocene Baltic amber, imaged using phase contrast X-ray computed micro-tomography. *Zootaxa* **2742**, 60-66.
- Penney, D., Green, D., Mcneil, A., Bradley, R., Marusik, Y., Withers, P., & Preziosi, R. 2012a. A new species of *Craspedisia* (Araneae: Theridiidae) in Miocene Dominican amber, imaged using X-ray computed tomography. *Paleontological Journal* **46**, 583-588.
- Penney, D., Mcneil, A., Green, D. I., Bradley, R. S., Jepson, J. E., Withers, P. J., & Preziosi, R. F. 2012b. Ancient Ephemeroptera–Collembola symbiosis fossilized in amber predicts contemporary phoretic associations. *PloS one* **7**, e47651
- Penney, D., Mcneil, A., Green, D. I., Bradley, R., Withers, P. J., & Preziosi, R. F. 2014. The oldest fossil pirate spider (Araneae: Mimetidae), in uppermost Eocene Indian amber, imaged using X-ray computed tomography. *Arachnology* **15**, 299-302.
- Peris, D., Davis, S. R., Engel, M. S. & Delclòs, X. 2014a. An evolutionary history embedded in amber: reflection of the Mesozoic shift in weevil-dominated (Coleoptera: Curculionoidea) faunas. *Zoological Journal of the Linnean Society* **171**, 534-553.
- Peris, D., Delclòs, X., Soriano, C., & Perrichot, V. 2014b. The earliest occurrence and remarkable stasis of the family Bostrichidae (Coleoptera: Polyphaga) in Cretaceous Charentes amber. *Palaeontologia Electronica* **17**, 1-8.
- Peris, D., Thayer, M. & Néraudeau, D. 2014c. Oldest Omaliini (Coleoptera: Staphylinidae: Omaliinae) Discovered in the Opaque Cretaceous Amber of Charentes. *Annals of the Entomological Society of America* **107**, 902-910.
- Peris, D., Philips, T. K., & Delclòs, X. 2015. Ptinid beetles from the Cretaceous gymnosperm-dominated forests. *Cretaceous Research* **52**, 440-452
- Perkovsky, E. E., Zosimovich, V. Y. & Vlaskin, A. P.. 2010. Rovno amber. In Penney, D. (eds) *Biodiversity of fossils in amber from the major world deposits*, 116-136. Manchester, UK: Siri Scientific Press.
- Perreau, M. 2012. Description of a new genus and two new species of Leiodidae (Coleoptera) from Baltic amber using phase contrast synchrotron X-ray microtomography. *Zootaxa* **3455**, 81-88.
- Perreau, M., & Perkovsky, E. E. 2014. Further description of *Catops nathani* Perkovsky 2001 from Late Eocene Baltic amber (Coleoptera: Leiodidae: Cholevinae: Cholevini) using phase contrast synchrotron X-ray microtomography. *Annales de la Société entomologique de France* **50**, 414-417.
- Perreau, M. & Tafforeau, P. 2011. Virtual dissection using phase-contrast X-ray synchrotron microtomography: reducing the gap between fossils and extant species. *Systematic Entomology*, **36**, 573-580.
- Perrichot, V. 2009. Long-tailed wasps (Hymenoptera: Megalyridae) from Cretaceous and Paleogene European amber. *Paleontological Contributions* **1**, 1-19.

- Perrichot, V., Nel, A. & Néraudeau, D. 2004. Two new wedge-shaped beetles in Albo-Cenomanian ambers of France (Coleoptera: Ripiphoridae: Ripiphorinae). *European Journal of Entomology* **101**, 583-589.
- Perrichot, V., Engel, M. S., Nel, A., Tafforeau, P. & Soriano, C. 2011. New earwig nymphs (Dermaptera: Pygidicranidae) in mid-Cretaceous amber from France. *Cretaceous Research* **32**, 325-330.
- Pike, E. M. 1994. Historical changes in insect community structure as indicated by hexapods of Upper Cretaceous Alberta (Grassy Lake) amber. *The Canadian Entomologist* **126**, 695-702.
- Pohl, H., Wipfler, B., Grimaldi, D., Beckmann, F., & Beutel, R. G. 2010. Reconstructing the anatomy of the 42-million-year-old fossil †*Mengeatertiaria* (Insecta, Strepsiptera). *Naturwissenschaften* **97**, 855-859.
- Poinar, G. O. 1992. *Life in amber*. Stanford University Press.
- Poinar, G. O. Jr. 1993. Amplification and sequencing of DNA from a 120–135-million-year-old weevil. *Nature* **363**, 536-538.
- Poinar, G. O., Zavortinik, T., Pike, T. & Johnston, P. 2000. *Paleoculicis minutus* (Diptera: Culicidae) n. Gen., n. Sp., from Cretaceous Canadian amber, with a summary of described fossil mosquitoes. *Acta Geologica Hispanica* **35**, 119-130.
- Poinar, G. & Danforth, B. 2006. A fossil bee from Early Cretaceous Burmese amber. *Science* **314**, 614-614.
- Quinney, A., Mays, C., Stilwell, J. D., Zelenitsky, D. K. & Therrien, F. 2015. The Range of Bioinclusions and Pseudoinclusions Preserved in a New Turonian (~90 Ma) Amber Occurrence from Southern Australia. *PLoS one* **10.5**, e0121307.
- Ragazzi, E. 2008. Fossil resin deposits or finds in Italy. *AMBERIF* **2008**, 5-40.
- Riedel, A., Dos Santos Rolo, T., Cecilia, A., Van De Kamp, T. 2012. Sayrevilleinae Legalov, a newly recognised subfamily of fossil weevils (Coleoptera, Curculionoidea, Attelabidae) and the use of synchrotron microtomography to examine inclusions in amber. *Zoological Journal of the Linnean Society* **165**, 773-794.
- Rowe, T., Cifelli, R. L., Lehman, T. M. & Weil, A. 1992. The Campanian Terlingua local fauna, with a summary of other vertebrates from the Aguja Formation, Trans-Pecos Texas. *Journal of Vertebrate Paleontology* **12**, 472-493.
- Saint Martin, S., Saint Martin, J-P., Girard, V. & Néraudeau, D. 2013. Organismes filamenteux de l'ambre du Santonien de Belcodene (Bouches-du-Rhône, France). *Annales de Paléontologie* **99**, 339-360.
- Saunders, W., Mapes, R., Carpenter, F. & Elsik, W. 1974. Fossiliferous amber from the Eocene (Claiborne) of the Gulf coastal plain. *Geological Society of America Bulletin* **85**, 979-984.
- Saupe, E. E., Pérez-De La Fuente, R., Selden, P. A., Delclos, X., Tafforeau, P. & Soriano, C. 2012. New *Orchestina* Simon, 1882 (Araneae: Oonopidae) from Cretaceous ambers of Spain and France: first spiders described using phase-contrast X-ray synchrotron microtomography. *Palaeontology* **55**, 127-143.
- Selden, P. A. 2002. First British Mesozoic spider, from Cretaceous amber of the Isle of Wight, southern England. *Palaeontology* **45**, 973-983.
- Shcherbakov, D. 2007. Extinct four-winged precoccids and the ancestry of scale insects and aphids (Hemiptera). *Russian Entomological Journal* **16**, 47-62.
- Shmakov, A., Perkovsky, E. 2009. Thrips (Thysanoptera, Insecta) from the Rovno amber, Eocene of Ukraine. *Paleontological Journal* **43**, 669-674.
- Sinitshenkova, N. D. 2000. A review of Triassic mayflies, with a description of new species from western Siberia and Ukraine (Ephemera = Ephemeroptera). *Paleontological Journal C/C OF Paleontologicheskii Zhurnal* **34**, S275-S283.

- Solórzano Kraemer, M. M., Perrichot, V., Brown, B. V., Tafforeau, P. & Soriano, C. 2011. A new species of the Cretaceous genus *Prioriphora* (Diptera: Phoridae) in French amber. *Systematic Entomology* **36**, 581-588.
- Solorzano Kraemer, M. M., Perrichot, V., Soriano, C. & Damgaard, J. 2014. Fossil water striders in Cretaceous French amber (Heteroptera: Gerromorpha: Mesoveliidae and Veliidae). *Systematic Entomology* **39**, 590-605.
- Sonibare, O. O., Hoffmann, T. & Foley, S. F. 2012. Molecular composition and chemotaxonomic aspects of Eocene amber from the Ameki Formation, Nigeria. *Organic geochemistry* **51**, 55-62.
- Soriano, C., Archer, M., Azar, D., Creaser, P., Delclòs, X., Godthelp, H., Hand, S., Jones, A., Nel, A. & Néraudeau, D. 2010. Synchrotron X-ray imaging of inclusions in amber. *Comptes Rendus Palevol* **9**, 361-368.
- Soriano, C., Pollock, D., Néraudeau, D., Nel, A. & Tafforeau, P. 2013. First fossil record of polypore fungus beetles from Lower Cretaceous amber of France. *Acta Palaeontologica Polonica* **59**, 941-946.
- Stankiewicz, B. A., Poinar, H. N., Briggs, D. E., Evershed, R. P. & Poinar, G. O. 1998. Chemical preservation of plants and insects in natural resins. *Proceedings of the Royal Society of London B: Biological Sciences* **265**, 641-647.
- Stroinski, A. & Szwedo, J. 2002. An overview of Fulgoromorpha and Cicadomorpha in East African copal (Hemiptera). *Denisia 04, Neue Folge* **176**, 57-66.
- Szadziewski, R. 1990. Biting midges (Insecta: Diptera: Ceratopogonidae) from Sakhalin amber. *Prace Muzeum Ziemi* **41**, 77-81.
- Szadziewski, R., Gilka, W. & Urbanek, A. 2015. A blood sucking biting midge from Upper Cretaceous Burmese amber with a key to the determination of fossil species in the relictual genus *Leptoconops* Skuse (Diptera: Ceratopogonidae). *Cretaceous Research* **54**, 255-259.
- Szwedo, J. 2002. Amber and amber inclusions of planthoppers, leafhoppers and their relatives (Hemiptera, Archeorrhyncha et Clypaeorrhyncha). *Denisia* **4**, 37-56.
- Van, T., Kamp, T., Rolo, S. & Baumbach, T. 2014. Scanning the Past—Synchrotron X-Ray Microtomography of Fossil Wasps in Amber. *Entomologie heute* **26**, 151-160.
- Vršanský, P. 2009. Albian cockroaches (Insecta, Blattida) from French amber of Archingeay. *Geodiversitas* **31**, 73-98.
- Waggoner, B. M. 1996. Bacteria and protists from Middle Cretaceous amber of Ellsworth County, Kansas. *PaleoBios* **17**, 20-26.
- Wang, B., Zhang, H. & Azar, D. 2011. The first Psychodidae (Insecta: Diptera) from the Lower Eocene Fushun amber of China. *Journal of Paleontology* **85**, 1154-1159.
- Ward, P. S. 1992. Ants of the genus *Pseudomyrmex* (Hymenoptera: Formicidae) from Dominican amber, with a synopsis of the extant Antillean species. *Psyche* **99**, 55-85.
- Wedmann, S., Makarkin, V. N., Weiterschan, T. & Hörnschemeyer T. 2013. First fossil larvae of Berothidae (Neuroptera) from Baltic amber, with notes on the biology and termitophily of the family. *Zootaxa* **3716**, 236-258.
- Weitschat, W. & Wichard, W. 2010. Baltic amber. In Penney, D. (ed.) *Biodiversity of fossils in amber from the major world deposits*, 80-115. Manchester, UK: Siri Scientific Press.
- Wimmer, R., Krumbiegel, G., Kosmowska-Ceranowicz, B. & Wagner-Wysiecka, E. 2015. New fossil resin discoveries in the Central German lignite mining area. *AMBERIF* **2015**, 8-30.