

Secular Trends of Birth Weight in Twins and Singletons in South Korea from 2000 to 2020

Yoon-Mi Hur

General College of Education, Kookmin University, Seoul, South Korea

Supplementary Table 1. The prevalence (%) of macrosomia and LBW, and preterm and term LBW in singletons and the first- and the second-born twins between 2000 and 2020.

		2000	2005	2010	2015	2020
Macrosomia	singletons	3.7	3.7	3.2	3.0	2.5
LBW	singletons	3.0	3.3	3.5	3.8	4.1
	1st-born twins	46.1	46.1	49.6	52.9	54.0
	2nd-born twins	51.9	53.8	58.1	59.5	61.4
LBW (preterm)	singletons	48.7	54.4	55.9	59.0	60.6
	1st-born twins	59.9	65.6	72.0	74.7	78.1
	2nd-born twins	55.2	61.2	67.6	70.9	75.0
LBW (term)	singletons	51.3	45.6	44.1	41	39.4
	1st-born twins	40.1	34.4	28.0	25.3	21.9
	2nd-born twins	44.8	38.8	32.4	29.1	25.0

Note. Macrosomia = birthweight > 4 kg; LBW = birthweight < 2.5 kg; LBW (term) = birthweight < 2.5 kg and gestational age \geq 37 weeks; LBW (preterm) = birthweight < 2.5 kg and gestational age < 37 weeks.