	Clinical			
Decision	Information	Delivery System	Health	Community
Support	System	Design	System	Resources
 Follow NCCN Guidelines on Smoking Cessation. Make 2A+R and Follow Up a routine, directive recommendation (i.e. Automatically refer patients to Quitline). Integrate specialist expertise to assist patients in smoking cessation. Provide culturally appropriate educational materials on the benefits of quitting smoking during treatment based on the NCCN Guidelines. 	 Build EHR Document tobacco status in system. Document acceptance of smoking cessation referral in system. Create alerts in system for providers on tobacco status of patients. Document discussion with and plan of care for tobacco users. Regularly collect and monitor process and outcome measures within 	 Train all providers in 2A+R and Follow Up. <u>www.helppatientsquitsc.org</u> Distribute tasks among team members who will be playing a role in assisting tobacco users to quit use. Offer internal cessation services. Develop internal and external processes (i.e. TelASK) for regular follow up with tobacco users and document status. 	 Adopt a policy within the practice/standard of care to follow the NCCN Guidelines on Smoking Cessation. Adopt and/or develop protocols for physicians and staff for providing smoking cessation therapies, both pharmacologic therapy and counseling. AAFP Pharma Chart Practice leadership establishes and monitors goals for decreasing tobacco use rates among cancer patients. Offer provider incentives based on quality of care. 	 Link patients to available loca community resources to assiss them in tobacco cessation. Form partnerships with organizations to support tobacco cessation. DHEC SC Tobacco Free Collaborative

- 100% of new cancer patients in the practice has their tobacco use and status documented in the patient record
- 100% of tobacco users within target populations with referrals to SC Tobacco Quitline (or other program) documented in the patient record
- 80% of tobacco users who do not initially enroll with the Quitline (or other program) who receive additional referrals and advise to quit tobacco use
- 80% of tobacco users in which Follow Up by provider on tobacco status is documented in the patient record