

**S1. Basic statistics of the population receiving services through our intervention  
(Acute phase)**

	Support status		p value
	Continuing (n=86) n (%) / maen (SD)	Terminated (n=39) n (%) / maen (SD)	
<b>Gender</b>			
Male	38 (44.2)	21 (53.8)	0.32*
Female	48 (55.8)	18 (46.2)	( $\chi^2=1.00$ )
<b>Age</b>			
19 years old and younger	1 (1.2)	2 (5.1)	
20-39 years old	8 (9.3)	7 (17.9)	
40-59 years old	32 (37.2)	5 (12.8)	<b><i>0.02*</i></b>
60 years old and older	45 (52.3)	25 (64.1)	<b><i>(<math>\chi^2=9.49</math>)</i></b>
<b>Living with family</b>			
Living with other family members	65 (75.6)	18 (46.2)	
Living alone	21 (24.4)	21 (53.8)	<b><i>&lt; 0.01*</i></b>
Unknown	0 (0.0)	0 (0.0)	<b><i>(<math>\chi^2=10.4</math>)</i></b>
<b>Job status</b>			
Employed/ Self-employed	12 (14.0)	5 (12.8)	
Unemployed	56 (65.1)	13 (33.3)	<b><i>&lt; 0.01*</i></b>
Unknown	18 (20.9)	21 (53.8)	<b><i>(<math>\chi^2=14.3</math>)</i></b>
Number of stress symptoms	2.42 (1.82)	1.39 (1.27)	<b><i>&lt; 0.01**</i></b> <b><i>(t=2.89)</i></b>

\*  $\chi^2$ -test, \*\* t-test

Excluded cases: age-unknown

***Italic*** : Significant variables ( $p < 0.05$ )

## S2. Association between support status and psychological/ social variables (Acute phase)

		Support status				Model 1		Model 2			
		Continuing (n=86)		Terminated (n=33)		Odds ratio (95% CI)	p value	Odds ratio (95% CI)	p value		
		n (%)	mean (SD)	n (%)	mean (SD)						
Past history of illness											
Physical illness	Yes	32	(37.2)	11	(33.3)	1.26	(0.53-2.96)	0.60	0.93	(0.38-2.32)	0.88
	No	54	(62.8)	22	(66.7)	1.00			1.00		
Mental illness	Yes	27	(31.4)	8	(24.2)	1.42	(0.56-3.56)	0.46	1.53	(0.58-4.02)	0.39
	No	59	(68.6)	25	(75.8)	1.00			1.00		
Disaster-related damage											
<u>Loss of family or relatives</u>	Yes	25	(29.1)	2	(6.1)	<b>6.51</b>	<b>(1.44-29.6)</b>	<b>&lt;0.01</b>	<b>5.79</b>	<b>(1.25-26.7)</b>	<b>0.02</b>
	No	61	(70.9)	31	(93.9)	1.00			1.00		
<u>Large-scale partial/ whole collapsed</u>	Yes	50	(58.1)	12	(36.4)	<b>2.38</b>	<b>(1.03-5.47)</b>	<b>0.04</b>	1.94	(0.82-4.62)	0.13
	No	36	(41.9)	21	(63.6)	1.00			1.00		
Current job											
	Employed/Self-employed	12	(17.6)	5	(15.2)	0.95	(0.30-3.05)	0.93	0.65	(0.18-2.27)	0.49
	Unemployment	56	(82.4)	28	(84.8)	1.00			1.00		
Economic & Resettlement issue											
	Yes	17	(19.8)	4	(12.1)	1.72	(0.53-5.58)	0.37	1.07	(0.30-3.85)	0.92
	No	69	(80.2)	29	(87.9)	1.00			1.00		
Alcohol-related problem											
	Yes	7	(8.1)	4	(12.1)	0.72	(0.19-2.75)	0.63	0.54	(0.13-2.18)	0.38
	No	79	(91.9)	29	(87.9)	1.00			1.00		
Isolation from family or community											
	Yes	10	(11.6)	6	(18.2)	0.69	(0.21-2.23)	0.53	0.70	(0.20-2.42)	0.57
	No	76	(88.4)	27	(81.8)	1.00			1.00		
<u>Number of stress symptoms</u>											
		2.42	(1.82)	1.39	(1.27)	<b>1.59</b>	<b>(1.14-2.22)</b>	<b>&lt;0.01</b>	-	-	-
Number of improved stress symptoms											
		1.27	(1.38)	1.15	(1.20)	1.08	(0.79-1.49)	0.63	0.57	(0.29-1.09)	0.09
<u>Support frequency (time/year)</u>											
		19.5	(20.9)	98.5	(73.8)	<b>0.95</b>	<b>(0.93-0.97)</b>	<b>&lt;0.01</b>	<b>0.95</b>	<b>(0.92-0.97)</b>	<b>&lt;0.01</b>
Rate of contact failure (%)											
		0.00	0.00	0.00	0.00	-	-	-	-	-	-

**Model 1:** independent variables were adjusted by sex and age

**Model 2:** independent variables were adjusted by sex, age and number of stress symptoms

**Excluded cases:** age-unknown, referred to psychiatric hospitals or clinics

**SD:** Standard Deviation **CI:** Confidence interval **Italic:** p<0.05