

Table S4. Odds ratios (95% CIs) from logistic regression models for maternal HDP and child motor or mental development abnormalities at the age of 3 years, stratified by SGA.

| Variables | No. of cases | No. of cases with HDP | | model 1 ^a | | model 2 ^b | | model 3 ^c | |
|--------------------|--------------|-----------------------|-------|----------------------|---------|----------------------|---------|----------------------|---------|
| | n | n | (%) | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Motor abnormality | | | | | | | | | |
| SGA + | 50 | 3 | (6.0) | 1.06 (0.33-3.43) | 0.927 | 0.62 (0.15-2.46) | 0.494 | 1.00 (0.31-3.29) | 0.993 |
| SGA - | 374 | 16 | (4.3) | 1.90 (1.15-3.15) | 0.013 | 1.45 (0.86-2.44) | 0.160 | 1.78 (1.07-2.96) | 0.026 |
| Mental abnormality | | | | | | | | | |
| SGA + | 54 | 1 | (1.9) | 0.31 (0.04-2.24) | 0.245 | 0.34 (0.05-2.58) | 0.298 | 0.27 (0.04-1.99) | 0.200 |
| SGA - | 493 | 26 | (5.3) | 2.39 (1.60-3.56) | <0.001 | 2.20 (1.46-3.30) | <0.001 | 2.14 (1.43-3.21) | <0.001 |

The reference group was defined as children born from mother without HDP in each group.

^aUnadjusted model.

^bMultivariable logistic models were adjusted for maternal age, multiple pregnancy, parity, maternal smoking, maternal drinking, sex of the child, gestational week, asphyxia, and jaundice.

^cMultivariable logistic models were adjusted for maternal age, multiple pregnancy, parity, maternal smoking, maternal drinking, sex of the child. OR, odds ratio; CI, confidence interval; HDP, hypertensive disorders of pregnancy; SGA, small for gestational age.