

2 **Amycenone reduces excess body weight and attenuates**  
3 **hyperlipidemia by inhibiting lipogenesis and promoting**  
4 **lipolysis and fatty acid  $\beta$ -oxidation in KK- $A^y$  obese diabetic**  
5 **mice**

6  
7 **Shortened version of the title**

8 Amycenone reduces excess body weight in KK- $A^y$  mice

9  
10 Maya Kudo,<sup>#a</sup> Misa Hayashi,<sup>a</sup> Boju Sun,<sup>c,d</sup> Lili Wu,<sup>e</sup> Tonghua Liu,<sup>c,d</sup> Ming Gao<sup>\*a,b</sup>

11 *a School of Pharmaceutical Science, Mukogawa Women's University. 11-68 Koshien Kyuban-cho,*  
12 *Nishinomiya, Hyogo 663-8179, Japan*

13 *b Institute for Bioscience, Mukogawa Women's University. 11-68 Koshien Kyuban-cho, Nishinomiya,*  
14 *Hyogo 663-8179, Japan*

15 *c Second Clinical Medical College, Beijing University of Chinese Medicine. No. 11 North Third Ring*  
16 *East Road, Beijing 100029, China*

17 *d Key Laboratory of Health Cultivation of the Ministry of Education, Beijing University of Chinese*  
18 *Medicine. No. 11 North Third Ring East Road, Beijing 100029, China*

19 *e College of traditional Chinese Medicine. Beijing University of Chinese Medicine. No. 11 North*  
20 *Third Ring East Road, Beijing 100029, China*

21  
22 **Author contributions**

23 **\*Corresponding author: Ming Gao (MG), PhD**

24 Professor, Mukogawa Women's University. 11-68 Koshien Kyuban-cho, Nishinomiya, Hyogo 663-  
25 8179, Japan

26 Email address: gaoming@mukogawa-u.ac.jp

27 Telephone and fax number: +81-798-45-9983

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

**#First author: Maya Kudo (MK)**

Email: 1422006@mwu.jp

**Authors' email addresses**

MK: 1422006@mwu.jp

MH: hayasi27@mukogawa-u.ac.jp

BS: sunbucm@163.com

LW: gingnido\_566@163.com

TL: thliu@vip.163.com

MG: gaoming@mukogawa-u.ac.jp

**Supplement file legends and table**

**Fig. S1 The phosphorylation levels of AMPK in EAT, SAT, PAT, and MAT of KK-A<sup>y</sup> mice with normal water or under amycenone treatment (0.76 g/kg body weight/day) for 8 weeks.**

Amycenone decreased the phosphorylation levels of AMPK in PAT and MAT of KK-A<sup>y</sup> mice. (A, G) EAT, (B, H) SAT, (C-E, I-K) PAT and (F, L) MAT. White bars represent the control group and black bars represent the amycenone group. The data value is given as means ± S.E.M. (n = 9, 8, respectively), \**P* < 0.05, \*\**P* < 0.01 vs. control group.

**Fig. S2 The phosphorylation and expression levels of FAS, ACC, HSL, CaMKK, LKB1, PKA, and C/EBPβ in PAT of KK-A<sup>y</sup> mice with normal water or under amycenone treatment (0.76 g/kg body weight/day) for 8 weeks.**

Amycenone increased phosphorylation levels of ACC, HSL, and PKA in PAT of KK-A<sup>y</sup> mice. (A, H) FAS, (B, I) ACC, (C, J) HSL, (D, K) CaMKK, (E, L) LKB1, (F, M) PKA and (G, N) C/EBPβ. White bars represent the control group and black bars represent the amycenone group. The data value is given as means ± S.E.M. (n = 9, 8, respectively), \**P* < 0.05, \*\**P* < 0.01 vs. control group.

**Fig. S3 The phosphorylation and expression levels of HSL, ACC, CaMKK, LKB1, Sirt1, PKA, PPAR $\gamma$ , C/EBP $\alpha$ , and C/EBP $\beta$  in MAT of KK-A $^y$  mice with normal water or under amycenone treatment (0.76 g/kg body weight/day) for 8 weeks.**

Amycenone enhanced phosphorylation levels of ACC, CaMKK, and PKA and reduced the expression level of PPAR $\gamma$  in MAT of KK-A $^y$  mice. (A, J) HSL, (B, K) ACC, (C, L) CaMKK, (D, M) LKB1, (E, N) Sirt1, (F, O) PKA, (G, P) PPAR $\gamma$ , (H, Q) C/EBP $\alpha$  and (I, R) C/EBP $\beta$ . White bars represent the control group and black bars represent the amycenone group. The data value is given as means  $\pm$  S.E.M. (n = 9, 8, respectively), \* $P$  < 0.05, \*\* $P$  < 0.01 vs. control group.

**Fig. S4 Signaling related to the inhibition of body weight gain in PAT and MAT of KK-A $^y$  mice with normal water or under amycenone treatment (0.76 g/kg body weight/day) for 8 weeks (A), (B).**

**Table S1 mRNA expression levels of lipid metabolism-related genes in PAT of KK-A $^y$  mice with normal water or under amycenone treatment (0.76 g/kg body weight/day) for 8 weeks.**

Genes	Control (%)	Amycenone (%)
<i>Atgl</i>	100.0 $\pm$ 1.41	93.2 $\pm$ 1.58
<i>Mcad</i>	100.0 $\pm$ 2.36	114.2 $\pm$ 0.92
<i>ap2</i>	100.0 $\pm$ 0.43	153.0 $\pm$ 0.51
<i>Cpt1</i>	100.0 $\pm$ 1.10	1196.3 $\pm$ 0.29*
<i>Adipor1</i>	100.0 $\pm$ 1.66	109.2 $\pm$ 0.55
<i>Adipor2</i>	100.0 $\pm$ 0.31	144.1 $\pm$ 0.79

Amycenone increased the gene expression level of *Cpt1* in PAT of KK-A $^y$  mice. The table shows the genes expression levels of *Atgl*, *Mcad*, *ap2*, *Cpt1*, *Adipor1*, and *Adipor2*. The data value is given as means  $\pm$  S.E.M. (n = 9, 8, respectively), \* $P$  < 0.05 vs. control group.